

Anomalous spin transport in a two-channel-Kondo quantum dot device

Prashant Sharma

Laboratory of Atomic and Solid State Physics, Cornell University, Ithaca NY 14853

(Dated: February 2, 2008)

We study the response of a two-channel Kondo quantum dot device proposed by Y. Oreg and D. Goldhaber-Gordon [Phys. Rev. Lett. **90**, 136602 (2003)] to a spin-bias applied across one of its channels formed by Fermi liquid reservoirs weakly coupled to a spin-1/2 quantum dot. When the temperature $T < T_K$, the Kondo temperature of the device, the spin conductance depends on the Kondo coupling of the dot spin to the other channel in an anomalous manner. For isotropic Kondo couplings to the two channels the spin conductance is quantized for $T \ll T_K$ characterizing the two-channel Kondo fixed point. On the other hand, for anisotropic couplings a crossover energy scale $T_A \neq 0$ determines the temperature $T \ll T_A$ when the spin conductance vanishes indicating one-channel Kondo behavior.

The Kondo effect^{1,2} is a well understood problem in condensed matter physics, and its experimental realization^{3,4,5,6,7,8} in quantum dot systems allows for a detailed investigation of several of its interesting theoretical features^{9,10,11} using transport measurements. The low-bias transport properties of an odd-electron quantum dot weakly coupled to two Fermi liquid reservoirs have been well described by the physics near one-channel Kondo fixed point¹²: When charge fluctuations in the dot can be neglected, the odd-electron spin of the quantum dot hybridizes with a single electron channel formed by a linear combination of electrons from the two reservoirs. The physics of multi-channel Kondo has, however, not been observed in such systems, although there have been a few theoretical proposals for observing two-channel Kondo physics in a non-equilibrium situation^{13,14,15}: When a large source-drain bias introduces decoherence between electrons in different reservoirs, so that they can be treated as two independent channels interacting with the spin of the quantum dot. Perhaps the most promising realization of the two-channel Kondo behavior in equilibrium is provided by an experimental set-up in which the quantum dot is coupled to two electron channels which have sufficiently strong repulsive electron-electron interactions to inhibit transfer of electrons from one channel to another. The importance of electron-electron interactions in stabilizing the two-channel Kondo fixed point, in the case when the channels are two identical Luttinger liquids, was first discussed in Ref. 16, and later, in the context of a quantum dot embedded in a carbon nanotube, in Ref. 17. The recent experimental proposal by Oreg and Goldhaber-Gordon, Ref. 18, of a modified single-electron transistor to observe two-channel Kondo physics, is based on similar ideas in an experimentally realizable geometry: An odd-electron quantum dot connected to a Fermi liquid reservoir, and also to a larger quantum island (see Fig. 1). A sufficiently large charging energy E_c of the quantum island inhibits particle transfer to and from the Fermi liquid reservoir leading to two-channel Kondo behavior^{18,19} when $T \ll E_c$. This set-up allows for charge transport measurements²⁰ to be made using the Fermi liquid reservoirs of the device. Furthermore, as recent experiments reported in Ref. 21 have realized a "spin battery", the two-channel Kondo device can be probed by spin transport as well. The "spin battery", which works by using adiabatic pumping of a chaotic cavity, provides a source for spin-bias which is controlled by the pumping frequency²² and by spin-orbit interac-




FIG. 1: Single electron transistor device of Ref. [18]. The two channels that hybridize with the dot are: (i) quantum island, and (ii) a linear combination of the electron channels Ψ_l and Ψ_r of the two Fermi liquid reservoirs.

tion effects²³ in the cavity.

The two-channel Kondo model that describes the low energy physics in the above mentioned system has several distinctive *non-Fermi liquid* features: (i) a logarithmic temperature dependence of the specific heat and magnetic susceptibility, (ii) a non-zero ground state entropy of magnitude $\frac{1}{2} \ln 2$. A simple description that captures these features is given in the language of abelian bosonization^{24,25,26,27} by introducing new spinor excitations that are non-locally related to the conduction electrons. In the treatment of Ref. 24, the Hamiltonian at the fixed point is written in terms of C -spinor degrees of freedom and a local fermion $\hat{d}$ representing the spin impurity. The entropy of the ground state that develops below the Kondo temperature, is associated with a local real (Majorana) fermion $\hat{d} + \hat{d}^\dagger$ that decouples from the conduction sea at the two-channel Kondo fixed point. The C -spinor of the "spin-flavour" sector couples to the local Majorana fermion $i[\hat{d}^\dagger - \hat{d}]$ only through the linear combination $C_X + C_X^\dagger$. The strength of this coupling is determined by the Kondo temperature T_K . This effective coupling implies that the phase variable $\hat{\theta}_X$ (conjugate to the difference in the number of spins between the two channels $\hat{N}_X$) is fixed at the bottom of a cosine potential well with a depth T_K . As a result, when the coupling is sufficiently strong ($T_K \gg T$), eigenstates of the fixed point Hamiltonian contain phase coherent superpositions of states

with the same number of total spin but different numbers of spins in the two channels²⁷.

In this paper anomalous effects are shown in spin transport in the aforementioned quantum dot system, because of *non*-Fermi liquid physics near the two-channel Kondo fixed point. The predicted effects are: (i) A perfect spin conductance for isotropic Kondo coupling to the two channels, when the charge conductance is made negligible by asymmetrically coupling the right and left Fermi liquid reservoirs to the quantum dot. (ii) A sharp change from spin conducting to spin insulating behavior when the Kondo couplings are made anisotropic, *i.e.*, there exists an anisotropy-dependent energy scale T_A such that for $T \ll T_A$ the system has vanishing spin conductance. Based on the picture mentioned above, a heuristic understanding of this behavior may be arrived at as follows. When $T_A = 0$ and $\hat{\theta}_X$ is a good quantum operator, a spin-bias λ across one of the channels transforms the phase $\hat{\theta}_X \rightarrow \hat{\theta}_X + \lambda t/2$. In the phase representation the current operator is $\partial_t \hat{\theta}_X / 2\pi$, and we immediately obtain a quantized spin-conductance in units of $\hbar/4\pi$. In the presence of anisotropy in the Kondo couplings, the ground state of the system is qualitatively different: the strongly coupled channel forms a singlet with the dot-spin, and $\hat{\theta}_X$ is no longer a good quantum operator. Therefore, the system is a spin insulator for energy scales below T_A , determined by the channel anisotropy. In what follows, we derive these results.

The modified single electron transistor can be described in terms of electron channels in the two Fermi liquid reservoirs $\Psi_{l,r}(x)$ and in the large quantum island $\Psi_2(x)$ (see Fig. 1), that hybridize with the spin-1/2 quantum dot. As we are interested in the low-energy physics, we linearize the electron excitation spectrum, and using open boundary conditions at the site of the single electron transistor ($x = 0$) write the fields as chiral left-moving fermions. Only a linear combination $\psi_1(x) = \Psi_l(x) \sin \theta_0 + \Psi_r(x) \cos \theta_0$ couples to the spin of the quantum dot at $x = 0$, while the independent field $\psi_0(x) = \Psi_l(x) \cos \theta_0 - \Psi_r(x) \sin \theta_0$ decouples²⁰. The angle $\theta_0 = \tan^{-1}[t_l/t_r]$ is dependent on the (real) tunneling amplitudes to the left (t_l) and right (t_r) Fermi liquid reservoirs.

The low-energy behavior of the above mentioned device, when the charge fluctuations in the quantum island are neglected, is that of the two-channel Kondo problem^{18,19}. In terms of *chiral* (left moving) Dirac fermions $\psi_{1,2}(x)$ ^{16,27} we can write the Hamiltonian ($\hbar = 1$):

$$\mathcal{H} = \sum_{j=1,2} \sum_{\sigma=\uparrow,\downarrow} \int dx \psi_{j\sigma}^\dagger(x) i v \partial_x \psi_{j\sigma}(x) - \sum_{j=1,2} \sum_{\sigma\sigma'} J_j \vec{S} \cdot \psi_{j\sigma'}^\dagger(0) \frac{\vec{\tau}_{\sigma'\sigma}}{2} \psi_{j\sigma}(0). \quad (1)$$

Here, $\vec{S}$ represents the spin-1/2 degree of freedom of the quantum dot, $\vec{\tau}$ are the Pauli matrices, and v is the Fermi velocity. Applying a spin chemical potential difference λ between the

right and left Fermi liquid leads adds a term

$$H_\lambda = \lambda \sum_{\sigma=\uparrow,\downarrow} \frac{\tau_{\sigma\sigma}^3}{2} [N_{l\sigma} - N_{r\sigma}], \quad (2)$$

$$N_{l\sigma} = \int_{-L/2}^{L/2} dx \Psi_{l\sigma}(x) \Psi_{l\sigma}(x), \quad (3)$$

to the Hamiltonian $\mathcal{H}$ in Eq. (1). This can be written in terms of the operator ψ_1 that couples to the spin of the dot, and the free field ψ_0 , using the relation:

$$N_{l\sigma} - N_{r\sigma} = \cos 2\theta_0 [N_{1\sigma} - N_{0\sigma}] - \sin 2\theta_0 N_{01\sigma}, \quad (4)$$

$$N_{01\sigma} = \int dx [\psi_{0\sigma}^\dagger(x) \psi_{1\sigma}(x) + H.c.]. \quad (5)$$

Having chosen the spin quantization axis, we use abelian bosonization to calculate the spin current. The four chiral fermions $\psi_{j\sigma}(x)$ can be written in terms of chiral bosons $\phi_{j\sigma}$:

$$\psi_{j\sigma}(x) = \frac{\chi_{j\sigma}}{\sqrt{2\pi a_0}} e^{-i\hat{\theta}_{j\sigma}} e^{-i\hat{N}_{j\sigma} \frac{2\pi}{L} x} e^{-i\phi_{j\sigma}(x)}. \quad (6)$$

Here v/a_0 is the bandwidth, $\hat{N}_{j\sigma}$ counts the change in the number of electrons with spin σ in channel j with respect to a free electron ground state, $\hat{\theta}_{j\sigma}$ is its conjugate phase operator, $L/2$ is the length of each channel, and the cocycles $\chi_{j\sigma}$ are required to satisfy the relations: $\{\chi_{j\sigma}, \chi_{j'\sigma'}\} = 2\delta_{jj'}\delta_{\sigma\sigma'}$. It is clear from the form of $\mathcal{H}$ that it can be written entirely in the spin sector (involving only the bosonic fields $\phi_{j\uparrow} - \phi_{j\downarrow}$). The only term that couples spin fields to the charge fields ($\phi_{j\uparrow} + \phi_{j\downarrow}$) is the $N_{01\sigma}$ term in the external perturbation H_λ .

In the experimental set-up the parameter θ_0 can be made small. As a result it is convenient to write the spin current operator I_{sp} as a sum of two distinct contributions:

$$I_{sp} = \frac{1}{2} \frac{d}{dt} \tau_{\sigma\sigma}^3 [N_{l\sigma} - N_{r\sigma}] = I_{11} + I_{01} \quad (7)$$

$$I_{11} = \cos 2\theta_0 \frac{1}{2} \frac{d}{dt} \sum_{\sigma} \tau_{\sigma\sigma}^3 N_{1\sigma} \quad (8)$$

$$I_{01} = \sin 2\theta_0 \frac{1}{2} \frac{d}{dt} \sum_{\sigma} \tau_{\sigma\sigma}^3 [\psi_{0\sigma}^\dagger \psi_{1\sigma} + H.c.] \quad (9)$$

It is important to note that the two-channel Kondo Hamiltonian (1) has spin- and charge- sectors separated, and also that the field ψ_0 does not couple to the dot spin. Therefore, the spin current $\langle I_{sp} \rangle$ can only depend on even powers of $\sin 2\theta_0$. This is easy to see for $\langle I_{11} \rangle$, since I_{11} depends only on the spin sector the lowest order contribution is quadratic in $V \sin 2\theta_0$. For $\langle I_{01} \rangle$, the lowest order contribution is from N_{01} term of H_λ . This is apparently of order $V \sin^2 2\theta_0$, however, it can easily be shown to vanish²⁰. It follows then, that there is no contribution to the zero bias conductance upto terms of order $\sin^2 2\theta_0$. Therefore, if one arranges the experimental set-up such that $\sin \theta_0 \ll 1$, we can neglect these contributions altogether. As we show below, the remaining dominant contribution to the spin current shows scaling behavior with temperature and externally applied spin bias λ . The θ_0 independent

spin current is in marked contrast to the charge current (in response to a charge-bias) that has been shown to have a peak value of $2e^2/h \sin^2 2\theta_0$ ²⁰. In what follows, we shall calculate the spin current $\langle I_{\text{sp}} \rangle$ in response to the externally applied bias $H_\lambda = \sum_\sigma \lambda \frac{\tau_{\sigma\sigma}}{2} N_{1\sigma}$. We begin by introducing *charge* (c), *spin* (s), *flavour* (f), and *spin-flavour* (X) bosons:

$$\begin{aligned}\Phi_{c(s)} &= \frac{1}{2} \sum_{j=1,2} [\phi_{j\uparrow} \pm \phi_{j\downarrow}], \quad \Phi_f = \frac{1}{2} \sum_\beta [\phi_{1\beta} - \phi_{2\beta}], \\ \Phi_X &= \frac{1}{2} [\phi_{1\uparrow} - \phi_{1\downarrow} - \phi_{2\uparrow} + \phi_{2\downarrow}].\end{aligned}\quad (10)$$

Similar relations hold for defining the corresponding zero-mode number $\hat{N}_{c,s,f,X}$ and phase operators $\hat{\theta}_{c,s,f,X}$. The physics near the two-channel-Kondo fixed point is conveniently described in terms of a new basis of *C*-spinors^{24,25}:

$$C_\alpha(x) = \frac{\hat{\chi}_\alpha}{\sqrt{2\pi a_0}} e^{-i\hat{\theta}_\alpha} e^{-i\hat{N}_\alpha 2\pi x/L} e^{-i\Phi_\alpha(x)}, \quad (11)$$

where $\alpha = c, s, f, X$. Following Ref. [24], a unitary transform $U = \exp[iS^z \Phi_s(0)]$ yields the Hamiltonian $\mathcal{H}' = U\mathcal{H}U^\dagger$ which has spin (s) and spin-flavour (X) sectors decoupled from the charge (c) and flavour (f) sectors. The Hamiltonian $\mathcal{H}'$ can be written in terms of the *C*-spinors and the local fermion $\hat{d} = \chi_s e^{i\hat{\theta}_s} S^-$, where $\hat{\theta}_s$ is the phase conjugate to the total spin number operator $\hat{N}_s$,

$$\begin{aligned}\mathcal{H}' &= U\mathcal{H}U^\dagger \\ &= i \int dx v \left[C_X^\dagger(x) \partial_x C_X(x) + C_s(x) \partial_x C_s(x) \right] \\ &+ \left\{ \frac{J_1}{\sqrt{2\pi a_0}} \hat{d} C_X^\dagger(0) + \frac{J_2}{\sqrt{2\pi a_0}} \hat{d} C_X(0) + H.c. \right\} \\ &+ \frac{1}{2} [J_1 + J_2 - 4\pi v] \left[\hat{d}^\dagger \hat{d} - \frac{1}{2} \right] : C_s^\dagger(0) C_s(0) : \\ &+ \frac{1}{2} [J_1 - J_2] \left[\hat{d}^\dagger \hat{d} - \frac{1}{2} \right] : C_X^\dagger(0) C_X(0) : \quad (12)\end{aligned}$$

In the absence of channel anisotropy $J_1 = J_2$, the above two-channel Kondo system flows to a fixed point with the critical coupling $[J_1 + J_2]/2 = J^* = 2\pi v$,^{16,26} which gives the Kondo temperature $J^*/\sqrt{2\pi a} \equiv \sqrt{T_K}$, where v/a is the reduced bandwidth. We restrict our attention to the effective Hamiltonian $\mathcal{H}_{2\text{CK}}$ that contains all the relevant (in the renormalization group sense) operators near the two-channel Kondo fixed point²⁸. This can be conveniently written by introducing Majorana (real) fermions

$$\begin{aligned}\hat{a} &= \frac{\hat{d} - \hat{d}^\dagger}{i\sqrt{2}}, \quad \hat{b} = \frac{\hat{d} + \hat{d}^\dagger}{\sqrt{2}} \\ \eta_a &= \frac{[C_X(x) + C_X^\dagger(x)]}{\sqrt{2}}, \quad \eta_b = \frac{[C_X(x) - C_X^\dagger(x)]}{\sqrt{2}}\end{aligned}\quad (13)$$

The effective Hamiltonian:

$$\begin{aligned}\mathcal{H}_{2\text{CK}} &= \frac{1}{2} \int dx \sum_{j=a,b} \eta_j(x) [iv\partial_x] \eta_j(x) \\ &- i2\sqrt{T_K} \hat{a} \eta_a(0) - i2\sqrt{T_A} \hat{b} \eta_b(0), \quad (14)\end{aligned}$$

valid for energy scales below the Kondo temperature T_K , also contains the crossover energy scale associated with channel anisotropy T_A . This energy scale is related to the Kondo couplings in Eq. (1) as²⁰ $T_A \sim [2\pi v(J_1 - J_2)/(J_1 + J_2)^2] T_K$. The external spin-bias in the basis of *C*-spinors

$$H'_\lambda = UH_\lambda U^\dagger = \frac{\lambda}{2} N_X - \frac{\lambda}{2} S^z, \quad (15)$$

where $N_X = \int dx : C_X^\dagger(x) C_X(x) :$ and $S^z = -i\hat{a}\hat{b}$ in terms of the local Majorana fermions. The corresponding spin current operator is

$$I'_{\text{sp}} = UI_{\text{sp}}U^\dagger = \frac{1}{2} \frac{d}{dt} [N_X + i\hat{a}\hat{b}] \quad (16)$$

A transformation $C_X \rightarrow C_X e^{i\lambda(x-vt)/2}$ removes the first term of H'_λ , and corresponds to a change in the chemical potential of the C_X fermions. We will calculate the energy dependent phase shift of these fermions as they scatter off the local Majorana fermions. As these are left-movers we choose the incoming states to the right of the impurity ($x > 0$). The equations of motion of these fermions are:

$$\partial_t \eta_a(x) = v\partial_x \eta_a(x) + \delta(x)\sqrt{T_K} \hat{a} \quad (17a)$$

$$\partial_t \hat{a} = -\frac{1}{2} \lambda \hat{b} - 2\sqrt{T_K} \eta_a(0) \quad (17b)$$

$$\partial_t \hat{b} = +\frac{1}{2} \lambda \hat{a} + 2\sqrt{T_A} \eta_b(0) \quad (17c)$$

Integrating across the impurity gives us:

$$2\sqrt{T_K} \hat{a} = \eta_a(0^-) - \eta_a(0^+) \quad (18a)$$

$$2\sqrt{T_A} \hat{b} = \eta_b(0^+) - \eta_b(0^-) \quad (18b)$$

$$2\partial_t \hat{a} = -\lambda \hat{b} - 2\sqrt{T_K} [\eta_a(0^+) + \eta_a(0^-)] \quad (18c)$$

$$2\partial_t \hat{b} = \lambda \hat{a} + 2\sqrt{T_A} [\eta_b(0^+) + \eta_b(0^-)] \quad (18d)$$

Introducing the scattering matrix $\mathcal{S}$ allows us to write the outgoing fields near the impurity

$$\eta_{Y=a,b}(0^-) = \sum_{Y'=a,b} \sum_k \mathcal{S}_{YY'}(k) \eta_{Y',k} \quad (19)$$

where $\eta_{Y,k}$ is the k -th mode of the incoming field $\eta_Y(0^+)$. Using this *ansatz* in Eqs. (18) we obtain:

$$\mathcal{S}_{aa} = 1 + \frac{(e^{i\delta_a} - 1)}{\left[1 + \left(\frac{\lambda}{8\sqrt{T_K}} \right)^2 (e^{i\delta_a} - 1)(e^{i\delta_b} - 1)/T_A \right]} \quad (20)$$

$$\mathcal{S}_{ba} = \frac{\lambda}{2} \sqrt{\frac{T_A}{T_K}} \frac{(e^{i\delta_b} - 1)}{4T_A} \mathcal{S}_{aa}, \quad (21)$$

$$e^{i\delta_a} = \frac{ivk + 2T_K}{ivk - 2T_K}, \quad e^{i\delta_b} = \frac{ivk + 2T_A}{ivk - 2T_A}. \quad (22)$$

The scattering phase shift of the C_X fermion at the two-channel Kondo fixed point (when $T_A = 0 = \lambda$) is easily verified to be $\delta_a(k=0) = \pi$, $\delta_b = 0$, in accordance with Ref. [25]. The spin current (16) can be written in terms of the




FIG. 2: Differential spin conductance g_{sp} as a function of temperature for three different values of the channel anisotropy energy scale T_A . Two-channel Kondo behavior is evident for $T_A \ll T \ll T_K$, while one-channel Kondo behaviour for $T \ll T_A$.

incoming and outgoing fields at the impurity, using (18), and evaluated at finite bias and temperature:

$$\begin{aligned} \langle I_{\text{sp}} \rangle &= v \langle C_X^\dagger(0^+) C_X(0^+) - C_X^\dagger(0^-) C_X(0^-) \rangle \\ &= \frac{v}{4\pi} \int dk \text{Re} [\mathcal{S}_{aa}^* \mathcal{S}_{bb} - 1 - |\mathcal{S}_{ab}|^2] \\ &\quad \times [f(vk - \lambda/2) - f(vk + \lambda/2)]. \end{aligned} \quad (23)$$

Here $f(x)$ is the Fermi function, and the integral is limited by the upper momentum cut-off T_K/v . At the two-channel Kondo fixed point ($T_A = 0$) the current at low bias $\lambda \ll T_K$,

$$\begin{aligned} \langle I_{\text{sp}} \rangle &\equiv \frac{\lambda}{4\pi} g_{\text{sp}} = -\frac{\lambda}{4\pi} \int dk \sin^2 \left(\frac{\delta_a}{2} \right) \frac{df(vk)}{dk} \\ g_{\text{sp}} &= \frac{2T_K}{\pi T} \Psi' \left(\frac{1}{2} + \frac{2T_K}{\pi T} \right) \approx 1 - \frac{1}{3} \left(\frac{\pi T}{2T_K} \right)^2. \end{aligned} \quad (24)$$

Here $\Psi'(x)$ is the derivative of the psi function, and the last

expression is valid for $T \ll T_K$. The quantized spin conductance g_{sp} is the signature of the two-channel Kondo fixed point. Deviations from the fixed point, because of $T_A \neq 0$ and also because of an external magnetic field at the quantum dot site, decrease the zero-bias spin conductance: At $T = 0$ a non-zero T_A leads to a vanishing zero-bias conductance, while

$$\begin{aligned} g_{\text{sp}} &= - \int dk \sin^2 \left(\frac{\delta_a - \delta_b}{2} \right) \frac{df(vk)}{dk} \\ &\approx \left(1 - \frac{T_A^2}{T_K^2} \right) \zeta(2) \left(\frac{T}{T_A} \right)^2, \text{ for } T \ll T_A \ll T_K \end{aligned} \quad (25)$$

The full temperature dependence is plotted in Fig. 2.

The role of local magnetic field is similar to that of channel anisotropy. The zero-bias spin conductance in the presence of a local magnetic field with Zeeman energy λ_h is obtained by replacing the spin-bias $\lambda \rightarrow \lambda_h$ in the phase shifts $\delta_{a,b}$. The Zeeman energy corresponding to the crossover energy T_A is then found to be $\lambda_h \sim \sqrt{T_A T_K}$. Note that we have shown the zero temperature behavior of spin conductance to be identical to that of the impurity entropy²⁹. Therefore, the presence of a degenerate ground state at the two-channel Kondo fixed point implies perfect conductance. Also, note that the temperature dependence of the spin conductance in this device, for $T_A = 0$, is similar to that of charge conductance in a one-channel Kondo set-up¹², *i.e.*, in the absence of the quantum island.

In conclusion, we have calculated the spin current through the two-channel-Kondo quantum dot device of Ref. 18 for temperature and spin-bias smaller than the Kondo temperature T_K . We have shown that the spin conductance is simply related to the phase shift of the C_X spinor, and that the two-channel Kondo fixed point can be identified by tuning the device parameters to obtain perfect spin conductance. Deviations from the fixed point, because of channel anisotropy and external magnetic field at the quantum dot site, are shown to lead to a spin insulator.

This work was supported by the Packard foundation.

- ¹ A. C. Hewson, *The Kondo Problem to Heavy Fermions* (Cambridge University Press, 1993).
- ² Ph Nozieres and A. Blandin, *J. Phys. (Paris)* **41**, 193 (1980).
- ³ D. Goldhaber-Gordon, *et al.*, *Nature* **391**, 156 (1998).
- ⁴ S. M. Cronenwert, T. H. Oosterkamp, and L. P. Kouwenhoven, *Science* **540**, 281 (1998).
- ⁵ D. Goldhaber-Gordon, *et al.*, *Phys. Rev. Lett.* **81**, 5225 (1998).
- ⁶ W. G. van der Wiel *et al.*, *Science* **289**, 2105 (2000).
- ⁷ J. Nygard, D. H. Cobden, and P. E. Lindelof, *Nature* **408**, 342 (2000).
- ⁸ W. Liang, M. Bockrath, and H. Park, *Phys. Rev. Lett.* **88**, 126801 (2002).
- ⁹ M. Pustilnik and L. I. Glazman, *Phys. Rev. Lett.* **87**, 216601 (2001).
- ¹⁰ I. Affleck and P. Simon, *Phys. Rev. Lett.* **86**, 2854 (2001).
- ¹¹ A. Kaminski, Y. V. Nazarov, and L. I. Glazman, *Phys. Rev. Lett.*

- 83**, 384 (1999).
- ¹² A. Kaminski, Y. V. Nazarov, and L. I. Glazman, *Phys. Rev. B* **62**, 8154 (2000).
- ¹³ X. G. Wen, *cond-mat/9812431v2*.
- ¹⁴ P. Coleman, C. Hooley, and O. Parcollet, *Phys. Rev. Lett.* **86**, 4088 (2001).
- ¹⁵ A. Rosch, J. Kroha, and P. Wölfle, *Phys. Rev. Lett.* **87**, 156802 (2001).
- ¹⁶ M. Fabrizio and A. O. Gogolin, *Phys. Rev. B* **51**, 17827 (1995).
- ¹⁷ E. H. Kim, G. Sierra, and C. Kallin, *cond-mat/0202387*.
- ¹⁸ Y. Oreg and D. Goldhaber-Gordon, *Phys. Rev. Lett.* **90**, 136602 (2003).
- ¹⁹ S. Florens and A. Rosch, *cond-mat/0311219*.
- ²⁰ M. Pustilnik, L. Borda, L. I. Glazman, and J. von Delft, *cond-mat/0309646*.
- ²¹ S. K. Watson, R. M. Potok, C. M. Marcus, and V. Umansky, *Phys.*

- Rev. Lett. **91**, 258301 (2003).
- ²² E. R. Mucciolo, C. Chamon, and C. M. Marcus, Phys. Rev. Lett. **89**, 146802 (2002).
- ²³ P. Sharma and P. W. Brouwer, Phys. Rev. Lett. **91**, 166801 (2003).
- ²⁴ V. J. Emery and S. Kivelson, Phys. Rev. B **46**, 10812 (1992).
- ²⁵ J. M. Maldacena and A. W. W. Ludwig, Nucl. Phys. B **506**, 565 (1997).
- ²⁶ Jinwu Ye, Phys. Rev. Lett. **79**, 1385 (1997).
- ²⁷ G. Zarand and J. von Delft, Phys. Rev. B **61**, 6918 (2000).
- ²⁸ I. Affleck, A. W. W. Ludwig, H. B. Pang, and D. L. Cox, Phys. Rev. B **45**, 7918 (1992).
- ²⁹ M. Fabrizio, A. O. Gogolin, and Ph. Nozierés, Phys. Rev. Lett. **74**, 4503 (1995).